

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problems Mailbox.**

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 : <b>C07J 17/00, 75/00, A61K 31/705</b>		(11) International Publication Number: <b>WO 97/31933</b>
A1		(43) International Publication Date: 4 September 1997 (04.09.97)
(21) International Application Number: <b>PCT/KR96/00123</b>		(74) Agents: <b>CHOI, Kyu, Pal et al.</b> ; 824-20, Yeoksam-dong, Kangnam-ku, Seoul 135-080 (KR).
(22) International Filing Date: 29 July 1996 (29.07.96)		
(30) Priority Data: 1996/4879 27 February 1996 (27.02.96) KR		
(71) Applicant (for all designated States except US): <b>CHEIL JE DANG CO. [KR/KR]</b> ; 500, Namdangmunro 5-ga, Chung-ku, Seoul 100-095 (KR).		
(71)(72) Applicant and Inventor: <b>PARK, Man, Ki [KR/KR]</b> ; Ccachi Village #310-1502, 66, Kumi-dong, Bundang-ku, Sungnam-shi, Kyungki-do 463-500 (KR).		
(72) Inventors; and		(81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GR, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).
(75) Inventors/Applicants (for US only): <b>LEE, Seung, Ki [KR/KR]</b> ; Hanshin Hankang, Apartment #5-803, 34-13, Bampo-dong, Seocho-ku, Seoul 137-040 (KR). <b>PARK, Jeong, Hill [KR/KR]</b> ; Hansol Village, Apartment #301-208, Ilwonbon-dong, Kangnam-ku, Seoul 135-230 (KR). <b>KIM, Jong, Moon [KR/KR]</b> ; Chukong 2nd, Apartment #229-313, Chamshil 2-dong, Songpa-ku, Seoul 138-222 (KR). <b>LEE, Kwang, Youl [KR/KR]</b> ; 105-1, Jayang-dong, Dong-ku, Daejeon 300-100 (KR). <b>HAN, Sang, Beom [KR/KR]</b> ; 1577-24, Pongchun 4-dong, Kwanak-ku, Seoul 151-054 (KR).		


Published

With international search report.


(54) Title: **NOVEL GINSENG SAPONIN COMPOUNDS, PROCESS FOR PREPARATION THEREOF AND ANTI-TUMOR AGENT COMPRISING THE SAME AS AN ACTIVE COMPONENT**

## (57) Abstract

The present invention relates to novel ginseng saponin compounds having a potent anti-tumor activity, which are represented by formulae (I) and (II). The compounds (I) and (II) above are novel and can be produced by heating plants of ginseng genus for 0.5 to 20 hours at a high temperature of 110 to 180 °C, or can be synthesized by acetylating ginsenoside Rg<sub>3</sub> and  $\Delta^{20(22)}$ -ginsenoside Rg<sub>3</sub> which are known ginseng saponin compounds, respectively. The present invention also relates to an anti-tumor composition comprising these compounds (I) and/or (II) as an active ingredient.



(I)



(II)

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Cote d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SS	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

WO 97/31933

PCT/KR96/00123

1


NOVEL GINSENG SAPONIN COMPOUNDS, PROCESS FOR PREPARATION  
THEREOF AND ANTI-TUMOR AGENT COMPRISING THE SAME AS AN  
ACTIVE COMPONENT

5


## TECHNICAL FIELD

The present invention relates to novel ginseng saponin  
10 compounds having an anti-tumor activity. More specifically,  
the present invention relates to novel ginseng saponin  
compounds having a potent anti-tumor activity, which are  
represented by the following formulas (I) and (II).

15



25



The present invention also relates to a process for  
preparation thereof, and an anti-tumor composition com-  
prising the same as an active component.

35

WO 97/31933

PCT/KR96/00123

2

## BACKGROUND ART

The physiological and biochemical researches in the effect of ginseng, particularly red ginseng, have been generally conducted on the subject of saponin components which are contained in large quantities in ginseng and has been known as the main component exhibiting the pharmacological effect of ginseng. However, the minor saponin components contained in red ginseng merely in minute quantities have been studied by very few groups heretofore because they can be hardly separated from ginseng.

## 15 DISCLOSURE OF INVENTION

Thus, the present inventors have concentratedly studied to find a method for enhancing the pharmacological effect of ginseng by treating ginseng under specific conditions to increase the contents of specific components and further for separating the respective components so that the study of their pharmacological effect can be made. As a result of such studies, we have identified that when a ginseng is heat-treated for 0.5 to 20 hours at a high temperature of 110 to 180°C, the contents of effective components which are present in a minor amount in ginseng increase, and consequently a processed ginseng having an enhanced pharmacological effect compared with fresh ginseng, white ginseng or red ginseng is prepared. In the procedure to determine the pharmacological effect of the various components separated from the processed ginseng, the present inventors have found novel components which have never been disclosed heretofore, and subsequently identified the chemical structure, pharmacological effect and process for preparation thereof. Thus, we have completed the present invention.

WO 97/31933

PCT/KR96/00123


3

Therefore, the present invention relates to saponin compounds identified as novel active components contained in ginseng.


5

The first object of the present invention is to provide the novel saponin compounds represented by the following formulas (I) and (II). The configuration at  $\Delta^{20(22)}$  of (II) is *zusammen* or *entgegen*.

10



15



20

25

It is a further object of the present invention to provide the process for preparing the novel ginseng saponin compounds having the formulas (I) and (II) above.

30

Further, it is still another object of the present invention to provide an anti-tumor composition comprising the compound (I) and/or (II) as an active component.

35

WO 97/31933

PCT/KR96/00123

4

## BEST MODE FOR CARRYING OUT THE INVENTION

The compounds of formulas (I) and (II) according to  
5 the present invention can be prepared by an extraction  
from the processed product of plant of *Panax* genus or by a  
synthetic method using known ginsenoside components as a  
starting material.

10 First, according to the extraction method, roots or  
leaves of the plant of *Panax* genus, for example, *Panax*  
*ginseng*, *Panax notoginseng*, *Panax quinquefolium*, *Panax*  
*japonicus*, etc., or tissue cultures thereof, or extracts  
therefrom with water or lower alcohol are heated for 0.5  
15 to 20 hours at a temperature of 110 to 180°C. The proc-  
essed ginseng thus obtained is extracted with water, or a  
suitable organic solvent, for example, lower alcohols such  
as methanol, ethanol, etc., or a solvent mixture thereof,  
and then the extract is concentrated under reduced pres-  
20 sure, suspended in water and then extracted with a nonpo-  
lar organic solvent such as hexane, ether, dichlorometh-  
ane, chloroform, ethylacetate or a solvent mixture there-  
of. The remaining aqueous layer is extracted with a  
polar organic solvent such as butanol and then the extract  
25 is subjected to chromatography to obtain a fraction con-  
taining compounds (I) and (II). This fraction is crys-  
tallized from a suitable solvent system, for example a  
solvent mixture of water and lower alcohol, preferably a  
solvent mixture of water and methanol in a ratio of 1:1 by  
30 volume, to prepare the desired pure saponin compounds (I)  
and (II).

According to this method, during the procedure of  
heat-treatment of ginseng, a sugar moiety attached to the  
35 20th carbon of panaxadiol saponins present in ginseng such  
as ginsenosides Ra, Rb<sub>1</sub>, Rb<sub>2</sub>, Rc, Rd, etc. is removed and  
an acetyl group is introduced into the 6th position of the

WO 97/31933

PCT/KR96/00123

5


terminal glucose of the sugar moiety attached to 3rd carbon to produce the novel saponin compound (I). The saponin compound (I) can also be produced by removing the sugar moiety attached to the 20th carbon from ginsenosides  
5  $Rs_1$  and  $Rs_2$ . The saponin compound (II) is produced by removing the OH group attached to the 20th carbon and hydrogen at the 22th-position from the compound (I) through dehydration reaction. In this reaction, the stereochemical structure of the double bond at 20th-position  
10 tion can have cis or trans configuration.

In the extraction method according to the present invention as mentioned above, the contents of the desired compounds (I) and (II) can be more increased by repeatedly  
15 carrying out chromatography. In addition, if necessary, the order of the heat-treatment step and extraction step with organic solvent in this process can be inverted to obtain the same result.

20 According to the synthetic method of the present invention, the novel saponin compounds of formulas (I) and (II) can be obtained by acetylating the known ginsenoside compounds. Specifically, the compound of formula (I) can be produced by acetylating the known ginsenoside  $Rg_3$  of  
25 formula (III), and the compound of formula (II) can be prepared by acetylating the known  $\Delta^{20(22)}$ -ginsenoside  $Rg_3$  of formula (IV) which is formed by the dehydration reaction at 20th-position of ginsenoside  $Rg_3$ .

30

35






WO 97/31933

PCT/KR96/00123

6

5



10 Such acetylation reaction can be conducted using  
acetic anhydride ( $\text{Ac}_2\text{O}$ ) or acetyl chloride as an acetylating  
agent. In this case, the acetylating agent can be  
used in a molar ratio of 1:1-4, preferably 1:1.2-2 with  
respect to the compound (III) or (IV). It is appropriate  
15 to carry out this reaction at a temperature of  $-40^\circ\text{C}$  to  
 $20^\circ\text{C}$ , preferably  $-40^\circ\text{C}$  to  $0^\circ\text{C}$ , for 1 to 48 hours.

The novel saponin compound of formula (I) or (II) thus  
obtained can be further purified by a conventional work-  
20 ing-up method, for example, selective crystallization,  
column chromatography, etc.

The novel saponin compounds (I) and (II) prepared  
according to the process of the present invention as  
25 mentioned above, have a potent anti-tumor activity, and  
therefore can be effectively used as an agent for preven-  
tion or treatment of cancerous disease such as hepatoma,  
gastric cancer, leukemia, etc. Therefore, the present  
invention also relates to an anti-tumor composition com-  
30 prising as an active ingredient the compound (I) or (II)  
or the mixture thereof.

When the composition containing the compounds (I)  
and/or (II) of the present invention is applied for clini-  
35 cal purposes as an anti-tumor agent, it can be combined  
with pharmaceutically acceptable carriers to prepare  
various formulations conventionally used in the pharmaceu-

WO 97/31933

PCT/KR96/00123

7

tical field, for example, oral preparations such as tablets, capsules, troches, solutions, suspensions, etc.; injectable preparations such as injectable solutions or suspensions, or ready-to-use dried powder which can be applied after re-constituted with injectable distilled water before injection, etc.; or locally applicable preparations such as ointments, creams, solutions, etc.

The carriers which can be used in the composition of the present invention are conventional ones in the pharmaceutical field, for example, binders, lubricants, disintegrating agents, excipients, solubilizers, dispersing agents, stabilizers, suspending agents, coloring agents, flavors and the like in the case of oral preparations; preservatives, agents for painlessness, solubilizers, stabilizers and the like in the case of injectable preparations; bases, excipients, lubricants, preservatives and the like in the case of locally applicable preparations. The pharmaceutical preparations thus produced can be administered orally, or parenterally such as for example intravenously, peritoneally, subcutaneously, or can be topically applied. In addition, the oral preparations may be administered together with an antacid or in the form of an enteric-coated preparation which is formulated by covering the orally administrable solid preparation such as tablet with the enteric coatings, in order to prevent decomposition of the preparation by gastric acid when it is administered per orally.

Although the administration dosage to a human being of the novel saponin compounds (I) and (II) according to the present invention can be selected depending on the absorption, inactivating rate and excretion rate of the active component in the body, age, sex and condition of the subject patient, severity of the disorders to be treated and the like, it is generally administered to an adult in an amount of 5 to 500mg, preferably 10 to 200mg daily.

WO 97/31933

PCT/KR96/00123

8

Therefore, when the composition of the present invention is formulated into the dosage unit form, each of the dosage unit form can contain the compounds (I) and/or (II) in an amount of 5 to 500mg, preferably 10 to 200mg on the basis of the effective amount range as mentioned above. If necessary, the dosage unit form thus formulated can be administered using a specialized method according to the judgement of the specialist who arranges or observes the administration and the requirement of the individuals. The total daily dosage can also be divided into several portions and administered over several times, preferably 1 to 6 times.

The present invention is more specifically explained by the following examples and test examples. However, it should be understood that the present invention is not limited to those examples in any manner.

EXAMPLE 1 : Preparation of ginseng extract containing the compounds (I) and (II)

100g of fresh ginseng was introduced into a sealed container and then heated for 2 hours at 130°C. The obtained processed ginseng was extracted with 200ml of methanol to obtain the methanol extract and then methanol was removed from the extract by evaporation. The remaining residue was suspended in 100ml of water, extracted 3 times with 100ml of ether, and then the remaining aqueous layer was extracted 3 times with 100ml of butanol saturated with water to obtain the butanol extract containing saponins. This butanol extract was dried and then subjected to silica gel column chromatography (eluent; ethyl acetate/methanol/ water = 20:1:1). According to the result of TLC analysis (developing solvent: ethyl acetate/methanol/ water = 10:1:1) of the eluates, 30mg of a fraction containing 50% of the desired compound (I) having the Rf value of 0.25 and 25mg of a fraction con-

WO 97/31933

PCT/KR96/00123

9

taining 55% of the compound (II) having the Rf value of 0.27 were obtained, respectively.

5 EXAMPLE 2 : Preparation of ginseng extract containing the compounds (I) and (II)

10 10kg of dried root-hair ginseng was extracted by refluxing with 20L of methanol for 4 hours. The ginseng extract thus obtained was dried under reduced pressure.

15 The resulting ginseng extract in the form of a syrup was introduced into an autoclave and then heated for 4 hours at 120°C. The heat-treated ginseng extract was subjected to silica gel column chromatography according to the same method as Example 1 and then the eluates were applied for

20 TLC analysis (developing solvent: ethyl acetate/methanol/water = 10:1:1) to obtain 3g of a fraction containing 50% of the desired compound (I) having the Rf value of 0.25 and 2g of a fraction containing 60% of the compound (II) having the Rf value of 0.27, respectively.

Example 3 : Preparation of the compound (I)

25 1g of the fraction containing the compound (I) prepared in Example 2 above was subjected to silica gel column chromatography according to the same manner as Example 1 using the mixed solvent of ethyl acetate/methanol/water(20:1:1) as an eluent to obtain 400mg of a fraction containing 92% of the desired compound (I). The

30 fraction thus obtained was crystallized from the solvent mixture of methanol/water (1:1, v/v) to obtain 200mg of the desired compound (I).

The compound (I) thus obtained exhibits the following

35 physico-chemical characteristics :

Chemical name : 3 $\beta$ ,12 $\beta$ ,20 $\beta$ -trihydroxy-damar-24-ene-3-O- $\beta$ -

WO 97/31933

PCT/KR96/00123

10

D-6"-O-acetyl-glucopyranosyl-(1→2)-β-D-glucopyranoside

Mass spectrum (FAB<sup>+</sup>, m/z) : 827 ([M+H]<sup>+</sup>), 849 ([M+Na]<sup>+</sup>)

5 (FAB<sup>-</sup>, M/Z) : 825 ([M-H]<sup>-</sup>)

CNMR (δ ppm, pyridine-d<sub>5</sub>) : 16.1, 16.2, 16.7, 17.1, 17.4,  
18.3, 20.9, 22.4, 22.7, 25.5, 26.3, 26.8, 27.7,  
30.8, 30.9, 34.8, 35.4, 36.7, 39.1, 39.4, 39.8,  
10 49.2, 50.1, 50.2, 51.6, 56.2, 62.5, 64.5, 70.7,  
70.8, 71.1, 73.6, 75.0, 75.1, 77.6, 77.7, 78.2,  
83.9, 88.9, 104.6, 105.8, 125.8, 130.5, 171.0

Example 4 : Preparation of the compound (II)

15

1g of the fraction containing the compound (II) prepared in Example 2 above was subjected to silica gel column chromatography according to the same manner as Example 1 using the solvent mixture of ethyl acetate/  
20 methanol/water(20:1:1) as an eluent to obtain 300mg of a fraction containing 95% of the desired compound (II). The fraction thus obtained was crystallized from the solvent mixture of methanol/water (1:1, v/v) to obtain 150mg of the desired compound (II).

25

The compound (II) thus obtained exhibits the following physico-chemical characteristics :

Chemical name : 3β,12β-dihydroxy-damar-20(22),24-diene-3-  
30 O-β-D-6"-O-acetyl-glucopyranosyl-(1→2)-β-D-glucopyranoside

Mass spectrum (FAB<sup>+</sup>, m/z) : 809 ([M+H]<sup>+</sup>), 831 ([M+Na]<sup>+</sup>)

35 CNMR (δ ppm, pyridine-d<sub>5</sub>) : 13.1, 15.7, 15.9, 15.9, 17.1,  
17.8, 18.5, 20.9, 25.7, 26.8, 27.1, 27.4, 28.1,  
32.2, 32.6, 35.4, 37.1, 39.3, 39.8, 40.3, 50.8,

WO 97/31933

PCT/KR96/00123

11

50.9, 51.1, 51.2, 56.5, 62.9, 64.8, 70.9, 71.1,  
71.5, 75.4, 77.9, 78.1, 78.1, 78.6, 84.3, 89.2,  
104.9, 106.2, 123.5, 125.2, 131.0, 140.8, 171.0

5 Example 5 : Synthesis of the compound (I)

50mg of ginsenoside Rg<sub>3</sub> was dried under reduced pressure and 1ml of 2,4,6-collidine was added thereto and then stirred for 10 minutes at -40°C. 10μl of acetyl chloride  
10 was introduced thereinto and then the mixture was allowed to react for 3 hours. The reaction mixture was warmed slowly to room temperature and allowed to stand for 1 hour at room temperature. 1ml of methanol was added and the reaction solution was subjected to silica gel column  
15 chromatography according to the same manner as Example 1 to obtain 20mg of the desired compound (I).

Example 6 : Synthesis of compound (II)

20 50mg of  $\Delta^{20(22)}$ -ginsenoside Rg<sub>3</sub> was dried under reduced pressure and 1ml of 2,4,6-collidine was added thereto and then stirred for 10 minutes at -40°C. 10μl of acetyl chloride was introduced thereinto and then the mixture was allowed to react for 3 hours. The reaction  
25 mixture was warmed slowly to room temperature and allowed to stand for 1 hour at room temperature. 1ml of methanol was added and the reaction solution was subjected to silica gel column chromatography according to the same manner as Example 1 to obtain 25mg of the desired compound  
30 (II).

Test Example 1 : Anti-tumor activity of the compounds (I) and (II)

35 The anti-tumor activity of the novel saponin compounds of formulas (I) and (II) according to the present invention was determined by the method for measuring the incor-

WO 97/31933

PCT/KR96/00123

12

poration amount of  $^3\text{H}$  thymidine as described in the following.

13.8g of DMEM (Dulbecco's Modified Eagle's Medium, 5 manufactured by Gibco Co.) was dissolved in 1L of deionized water and then adjusted to pH 7.4 with sodium carbonate and hydrochloric acid solution. Then, 10% calf serum,  $1 \times 10^{-7}\text{M}$  of insulin and 50mg/L of gentamycin were added thereto. The mixture was then sterilized by means 10 of a millipore filter to prepare the culture solution. To this culture solution was inoculated human hepatoma sk-Hep-1 cell line, which was distributed from Cancer Research Center of Seoul National University in Korea, in a ratio of  $1 \times 10^6$  cells per  $25\text{cm}^2$  of the T flask area, which 15 was then incubated for 48 hours in an incubator of  $37^\circ\text{C}$  while keeping 5%  $\text{CO}_2$  gas. The culture product was transferred to a 24-well incubator and subcultured for one day, and then each of the compounds (I) and (II) dissolved in 70% ethanol was added thereto to a concentration of 0.01 20 to  $10\mu\text{M}$ , respectively. The same volume of 70% ethanol, instead of compounds (I) and (II), was added to the control group. 12 hours after treatment with each of the compounds (I) and (II),  $^3\text{H}$ -labelled thymidine was added to a concentration of  $1\mu\text{Ci/ml}$ . After 12 hours, the medium 25 was removed from each well and the cells were fixed with methanol, washed with PBS and then washed twice with 10% trichloroacetic acid to remove the unreacted radioactive thymidine. The cells were dissolved in 1N sodium hydroxide solution and neutralized with 1N hydrochloric acid and 30 then the radioactivity introduced into DNA was measured by scintillation counter (Pharmacia 1024). The measured results are described in the following Table 1.

35

WO 97/31933

PCT/KR96/00123

13

Table 1. Incorporation amount of radioactive thymidine into the DNA of hepatoma sk-Hep-1 cells depending on the concentration of compounds (I) and (II).

Concentration ( $\mu$ M)	Compound (I)		Compound (II)	
	dpm/well (Mean $\pm$ S.E.)	Percentage to control	dpm/well (Mean $\pm$ S.E.)	Percentage to control
Control	17812 $\pm$ 819	100.0	25828 $\pm$ 1918	100.0
0.01	23281 $\pm$ 1256	130.7	19242 $\pm$ 570	74.5
0.1	15016 $\pm$ 1231	84.3	15658 $\pm$ 1873	60.6
0.5	15665 $\pm$ 921	87.9	15137 $\pm$ 1467	58.6
1	15729 $\pm$ 302	88.3	13646 $\pm$ 2127	52.8
2.5	13155 $\pm$ 563	73.9	6421 $\pm$ 1527	24.9
5	733 $\pm$ 145	4.1	430 $\pm$ 105	1.7
10	728 $\pm$ 254	4.1	457 $\pm$ 115	1.8

Note S.E. : Standard error

As can be seen from the results described in Table 1 above, both the compounds (I) and (II) remarkably decrease the amount of incorporated radioactive thymidine at a concentration of 0.1 $\mu$ M or more, particularly 5 $\mu$ M or more. Therefore, it can be seen that the compounds (I) and (II) significantly inhibit the growth of hepatoma sk-Hep-1 cells.

Test Example 2 : Cell growth-inhibitory activity of the compounds (I) and (II) against human hepatoma cells

13.8g of DMEM (Dulbecco's Modified Eagle's Medium, manufactured by Gibco Co.) was dissolved in 1L of deionized water and then adjusted to pH 7.4 with sodium carbon-



WO 97/31933

PCT/KR96/00123

14

ate and hydrochloric acid solution. Then, 10% calf serum,  $1 \times 10^{-7} \text{M}$  of insulin and 50mg/L of gentamycin were added thereto. The mixture was then sterilized by means of a millipore filter to prepare the culture solution.

5 To this culture solution was inoculated human hepatoma cell line sk-Hep-1, which was distributed from Cancer Research Center of Seoul National University, in a ratio of  $1 \times 10^6$  cells per  $25 \text{cm}^2$  of the T flask area, which was then incubated for 48 hours in an incubator of  $37^\circ\text{C}$  while

10 keeping 5% of  $\text{CO}_2$  gas. The culture product was transferred to a 96-well incubator in a concentration of  $10^3$  cells per well and subcultured for one day, and then each of the compounds (I) and (II) dissolved in 70% ethanol was added thereto to a concentration of 0.1 to  $50 \mu\text{M}$ , respectively.

15 After 24 and 48 hours from the treatment with each of the compounds (I) and (II),  $20 \mu\text{l}$  of 3-[4,5-dimethyl-thiazol-2-yl]-2,5-diphenyl tetrazolium bromide (MTT) dissolved in PBS(phosphate buffered saline) (5mg/ml) was added thereto and treated for 4 hours at  $37^\circ\text{C}$  to produce

20 the insoluble formazane. The reaction mixture was centrifuged and the supernatant was removed. Then,  $100 \mu\text{l}$  of DMSO(dimethylsulfoxide) was added to dissolve the formazane precipitate and then the optical density at 570nm was measured by means of automatic plate reader as an index of

25 the amount of formazane thus produced. The measured results are described in the following Tables 2 (after 24 hours) and 3 (after 48 hours).

30

35

WO 97/31933

PCT/KR96/00123

15

Table 2. MTT analysis against human hepatoma sk-Hep-1 cells depending on the concentration of compounds (I) and (II) (after 24 hours)

Concentration ( $\mu$ M)	Compound (I)		Compound (II)	
	O.D. (Mean $\pm$ S.D.)	Percentage to control	O.D. (Mean $\pm$ S.D.)	Percentage to control
Control	0.98 $\pm$ 0.03	100.0	0.94 $\pm$ 0.16	100.0
0.01	0.88 $\pm$ 0.05	89.7	0.70 $\pm$ 0.02	74.6
0.1	0.74 $\pm$ 0.01	76.0	0.65 $\pm$ 0.00	69.5
0.5	0.68 $\pm$ 0.05	69.6	0.59 $\pm$ 0.04	62.7
1	0.61 $\pm$ 0.03	61.9	0.52 $\pm$ 0.03	55.9
5	0.50 $\pm$ 0.01	50.7	0.46 $\pm$ 0.04	48.7
10	0.42 $\pm$ 0.04	43.1	0.36 $\pm$ 0.01	38.2
25	0.17 $\pm$ 0.03	17.3	0.24 $\pm$ 0.03	26.0

Table 3. MTT analysis against human hepatoma sk-Hep-1 cells depending on the concentration of compounds (I) and (II) (after 48 hours)

Concentration ( $\mu$ M)	Compound (I)		Compound (II)	
	O.D. (Mean $\pm$ S.D.)	Percentage to control	O.D. (Mean $\pm$ S.D.)	Percentage to control
Control	1.02 $\pm$ 0.02	100.0	1.07 $\pm$ 0.04	100.0
0.01	0.89 $\pm$ 0.04	87.8	0.90 $\pm$ 0.04	83.8
0.1	0.70 $\pm$ 0.03	69.0	0.69 $\pm$ 0.03	64.1
0.5	0.42 $\pm$ 0.02	41.6	0.42 $\pm$ 0.01	38.7
1	0.34 $\pm$ 0.02	33.0	0.36 $\pm$ 0.01	33.8
5	0.18 $\pm$ 0.01	18.1	0.28 $\pm$ 0.04	26.0

WO 97/31933

PCT/KR96/00123

16

Table 3. (continued)

5	Compound (I)			Compound (II)	
	Concentration ( $\mu$ M)	O.D. (Mean $\pm$ S.D.)	Percentage to control	O.D. (Mean $\pm$ S.D.)	Percentage to control
10	10	0.14 $\pm$ 0.01	13.7	0.14 $\pm$ 0.01	12.7
	25	0.12 $\pm$ 0.01	11.6	0.12 $\pm$ 0.01	11.0

Note O.D. : Optical density  
S.D. : Standard deviation

15

From the results described in Tables 2 and 3 above, it can be seen that since the optical density(O.D.) decreases accordingly as the concentration of the compound (I) or (II) increases, the compound of the present invention significantly inhibits the cell-growth of hepatoma sk-Hep-1 cells.

Test Example 3 : Acute toxicity test of the compounds (I) and (II)

25

40 mice weighing 20 to 40g were used as test animal and divided into 2 groups including 20 mice, respectively. Each of the compounds (I) and (II) according to the present invention was suspended in 1ml of physiological saline and orally administered to each group. After 14 days from administration, the number of survived test animal was counted. To the control group, 1ml of physiological saline was orally administered. The results are described in the following Table 4.

35

WO 97/31933

PCT/KR96/00123

17

Table 4. Acute toxicity of the compounds (I) and (II)  
against mouse

5	Test group	Dosage (mg/kg, Oral)	Number of Test animals	Number of Survived animals
	A	1000	20	20
10	B	1000	20	20

Note A : Compound (I) receiving group  
B : Compound (II) receiving group

15

From the results described in Table 4 above, it can be  
seen that the novel ginseng saponin compounds (I) and (II)  
according to the present invention have no substantial  
20 toxicities.

25

30

35


WO 97/31933

PCT/KR96/00123


18

WHAT IS CLAIMED IS :

1. A ginseng saponin compound having the following formula (I):




- 15 2. A ginseng saponin compound having the following formula (II) wherein the configuration of  $\Delta^{20(22)}$  is zusammen or entgegen:



25

3. A process for preparing a compound having the following formula (I):

30




WO 97/31933

PCT/KR96/00123

19

which comprises acetylating ginsenoside Rg<sub>3</sub> having the following formula (III):


5



10

4. A process for preparing a compound having the following formula (II):


15



20

which comprises acetylating  $\Delta^{20(22)}$ -ginsenoside Rg<sub>3</sub> having the following formula (IV):

25



30

- 35 5. A process for preparing the compounds of formulas (I) and (II) as defined in claims 1 and 2, characterized in that an extract from the plant of Panax genus with

WO 97/31933

PCT/KR96/00123

20

water or lower alcohol is heated for 0.5 to 20 hours at a temperature of 110 to 180°C, the processed ginseng thus obtained is extracted with water, an organic solvent or a mixture thereof and then the extract is concentrated under reduced pressure, suspended in water and then extracted with a nonpolar organic solvent, the aqueous layer is separated and extracted with a polar organic solvent, and the obtained extract is subjected to chromatography to obtain a fraction containing the said compounds (I) and (II), which is then crystallized from a solvent mixture of water and lower alcohol.

6. An anti-tumor composition comprising the compound of formula (I) as defined in claim 1, the compound of formula (II) as defined in claim 2 or a mixture thereof as an active component, together with a pharmaceutically acceptable carrier.

7. The composition of claim 6 which is formulated into a pharmaceutical dosage unit form.

25

30

35

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/KR 96/00123

## A. CLASSIFICATION OF SUBJECT MATTER

IPC<sup>6</sup>: C 07 J 17/00, 75/00; A 61 K 31/705

According to International Patent Classification (IPC) or to be national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC<sup>6</sup>: C 07 J 17/00, 75/00; A 61 K 31/705

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

QUESTEL: G-DARC

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	Chemical Abstracts, Vol.112, No. 19, 07 May 1990 (Columbus, Ohio, USA), page 432, abstract No.175597h, I.KITAGAWA et al., "Chemical studies on crude drug processing. VI. Chemical structures of malonyl-ginsenosides Rb <sub>1</sub> , Rb <sub>2</sub> , Rc and Rd isolated from the root of Panax ginseng C.A. Meyer", & Chem.Pharm.Bull. 1989, 37(11) 2961-70.	1,5
A	Chemical Abstracts, Vol.123, No.9, 28 August 1995 (Columbus, Ohio, USA), page 748, abstract No. 122844g, D.S.KIM et al., "Preparation and structur determination of a new glycoside, (20E)-ginsenoside Rh <sub>3</sub> , and its isomer from diol-type ginseng saponins.", & Yakhak Hoechi 1995, 39(1) 85-93.	2,5
A	Chemical Abstracts, Vol.107, No.3, 20 July 1987 (Columbus, Ohio, USA), page 642, abstract No. 23596r, L.N.ATOPKINA et al., "Glycosylation of dammarane type triterpenoids. IV. β-D-Glucopyranosides of betulafolienetriol and its derivatives", & Khim. Priir. Soedin. 1968, (3), 301-12.	1

☒ Further documents are listed in the continuation of Box C☒ See patent family annex

## \* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, etc., exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

- "&" document member of the same patent family

Date of the actual completion of the international search

16 January 1997 (16.01.97)

Date of mailing of the international search report

24 January 1997 (24.01.97)

Name and mailing address of the ISA/AT  
AUSTRIAN PATENT OFFICE  
Kohlmarkt 8-10  
A-1014 Vienna  
Facsimile No. 1/53424/535

Authorized officer

Hofbauer

Telephone No.

1/53424/225

Form PCT/ISA/210 (second sheet) (July 1992)



## INTERNATIONAL SEARCH REPORT

International application No.

PCT/KR 96/00123

## C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	DE 40 01 895 A1 (HARRIER GMBH) 25 July 1991 (25.07.91), pages 1-7.	2,6,7

Form PCT/ISA/210 (continuation of second sheet) (July 1992)

